

MAT 1575 Final Exam Review Problems

Revised by Prof. Kostadinov Spring 2014, Prof. ElHitti Summer 2017, Prof. Africk Spring 2026

1. Evaluate the following definite integrals:

$$\text{a. } \int_0^1 x^2(x^3 + 1)^3 dx \quad \text{b. } \int_0^1 \frac{x}{\sqrt{x^2 + 9}} dx \quad \text{c. } \int_0^1 \frac{3x^2}{\sqrt[3]{x^3 + 1}} dx$$

2. Evaluate the following indefinite integrals:

$$\text{a. } \int x^2 \ln(x) dx \quad \text{b. } \int x^2 e^{-x} dx \quad \text{c. } \int x \cos(3x) dx$$

3. Find the area of the region enclosed by the graphs of:

$$\text{a. } y = 3 - x^2 \text{ and } y = -2x \quad \text{b. } y = x^2 - 2x \text{ and } y = x + 4$$

4. Find the volume of the solid obtained by rotating the region bounded by the graphs of:

$$\begin{aligned} \text{a. } y = x^2 - 9, y = 0 \text{ about the x-axis.} \quad & \text{b. } y = 16 - x, y = 3x + 12, x = -1 \text{ about the x-axis.} \\ \text{c. } y = x^2 + 2, y = -x^2 + 10, x \geq 0 \text{ about the y-axis.} \end{aligned}$$

5. Evaluate the following indefinite integrals:

$$\text{a. } \int \frac{1}{x^2 \sqrt{36 - x^2}} dx \quad \text{b. } \int \frac{\sqrt{x^2 - 9}}{x^4} dx \quad \text{c. } \int \frac{9}{x^2 \sqrt{x^2 + 9}} dx \quad \text{d. } \int \frac{6}{x^2 \sqrt{x^2 - 36}} dx$$

6. Evaluate the following indefinite integrals:

$$\text{a. } \int \frac{3x+7}{x^2+6x+9} dx \quad \text{b. } \int \frac{5x+6}{x^2-36} dx \quad \text{c. } \int \frac{3x+2}{x^2+2x-8} dx \quad \text{d. } \int \frac{12-8x}{x^2(x-6)} dx \quad \text{e. } \int \frac{-2x^2+4x+4}{x(x-2)^2} dx$$

7. Evaluate the improper integral:

$$\text{a. } \int_0^{\infty} \frac{2}{(x+2)^3} dx \quad \text{b. } \int_0^{\infty} \frac{5}{\sqrt[5]{x+5}} dx \quad \text{c. } \int_3^5 \frac{3}{\sqrt[3]{(x-3)^4}} dx$$

8. Decide if the following series converges or not. Justify your answer using an appropriate test:

$$\text{a. } \sum_{n=1}^{\infty} \frac{9n^5}{3n^5+5} \quad \text{b. } \sum_{n=1}^{\infty} \frac{5}{10^n} \quad \text{c. } \sum_{n=1}^{\infty} \frac{5n}{10^n} \quad \text{d. } \sum_{n=1}^{\infty} \frac{n!}{n^2 5^n} \quad \text{e. } \sum_{n=1}^{\infty} \left(\frac{n+1}{2n+3} \right)^n$$

9. Determine whether the series is absolutely or conditionally convergent or divergent:

$$\text{a. } \sum_{n=1}^{\infty} (-1)^n \frac{10}{7n+2} \quad \text{b. } \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n^5}} \quad \text{c. } \sum_{n=0}^{\infty} (-1)^n 5^{-n} \quad \text{d. } \sum_{n=1}^{\infty} (-1)^n \frac{n^2 - n - 1}{2n^2 + n + 1}$$

10. Find the radius and the interval of convergence of the following power series:

$$a. \sum_{n=0}^{\infty} \frac{(x-1)^n}{n+2} \quad b. \sum_{n=0}^{\infty} \frac{(-1)^n(x-1)^n}{n+2} \quad c. \sum_{n=1}^{\infty} \frac{(x+1)^n}{n5^n} \quad d. \sum_{n=1}^{\infty} \frac{(-1)^n(x+1)^n}{n5^n}$$

11. Find the Taylor polynomial of degree 2 for the given function, centered at the given number a :

$$a. f(x) = e^{-2x} \text{ at } a = -1. \quad b. f(x) = \cos(5x) \text{ at } a = 2\pi.$$


12. Find the Taylor polynomial of degree 3 for the given function, centered at the given number a :

$$a. f(x) = 1 + e^{-x} \text{ at } a = -1 \quad b. f(x) = \sin(x) \text{ at } a = \frac{\pi}{2}$$

Answers:


$$(1a). \frac{5}{4} \quad (1b). \sqrt{10} - 3 \quad (1c). \frac{3}{2} (2^{2/3} - 1)$$

$$(2a). \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} + C \quad (2b). -(x^2 + 2x + 2)e^{-x} + C \quad (2c). \frac{1}{3} x \sin(3x) + \frac{1}{9} \cos(3x) + C$$




(3a). The area of the region between the two curves is:

$$Area = \int_{-1}^3 (3 - x^2 - (-2x)) dx = \frac{32}{3}$$



(3b). The area of the region between the two curves is:

$$Area = \int_{-1}^4 (x + 4 - (x^2 - 2x)) dx = \frac{125}{6}$$



(4a). Approximate the volume of the solid by vertical disks with radius $y = x^2 - 9$ between $x = -3$ and $x = 3$; gives the volume is $V = \int_{-3}^3 \pi(x^2 - 9)^2 dx = \frac{1296}{5} \pi$.

(4b). Using a washer of outer radius $R_{outer} = 16 - x$ and inner radius $R_{inner} = 3x + 12$ at x , gives the volume:

$$V = \pi \int_{-1}^1 ((16 - x)^2 - (3x + 12)^2) dx = \frac{656\pi}{3}, \text{ where the}$$

upper limit 1 is obtained from $16 - x = 3x + 12 \Rightarrow x = 1$.

(4c). 16π

$$(5a). -\frac{\sqrt{36-x^2}}{36x} + C \quad (5b). \frac{(x^2-9)^{3/2}}{27x^3} + C \quad (5c). -\frac{\sqrt{x^2+9}}{x} + C \quad (5d). \frac{\sqrt{x^2-36}}{6x} + C$$

$$(6a). \frac{2}{x+3} + 3\ln|x+3| + C \quad (6b). 3\ln|x-6| + 2\ln|x+6| + C \quad (6c). \frac{5}{3}\ln|x+4| + \frac{4}{3}\ln|x-2| + C$$

$$(6d). \frac{2}{x} - \ln|x-6| + \ln|x| + C \quad (6e). \ln|x| - \frac{2}{x-2} - 3\ln|x-2| + C$$

$$(7a). \frac{1}{4} \quad (7b). \text{The integral does not converge} \quad (7c). \text{The integral does not converge}$$

$$(8a). \lim_{n \rightarrow \infty} \frac{9n^5}{3n^5+5} = \lim_{n \rightarrow \infty} \frac{3}{1+5/3n^5} = 3 > 0 \text{ so the series diverges by the test for divergence.}$$

(8b). This is a geometric series, with common ratio $r = 1/10 < 1$, so it converges to $5/9$:

$$\sum_{n=1}^{\infty} \frac{5}{10^n} = \frac{a}{1-r} = \frac{5/10}{1-\frac{1}{10}} = \frac{5/10}{9/10} = \frac{5}{9}$$

$$(8c). \lim_{n \rightarrow \infty} \frac{5(n+1)}{10^{n+1}} / \frac{5n}{10^n} = \lim_{n \rightarrow \infty} \frac{1+1/5n}{10} = \frac{1}{10} < 1 \text{ so the series converges by the ratio test.}$$

$$(8d). \lim_{n \rightarrow \infty} \frac{(n+1)!}{(n+1)^2 5^{n+1}} / \frac{n!}{n^2 5^n} = \lim_{n \rightarrow \infty} \frac{n^2}{5(n+1)} = \lim_{n \rightarrow \infty} \frac{n}{5(1+\frac{1}{n})} = \lim_{n \rightarrow \infty} \frac{n}{5} = \infty \text{ so the series diverges by the ratio test.}$$

$$(8e). \lim_{n \rightarrow \infty} \left[\left(\frac{n+1}{2n+3} \right)^n \right]^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n+1}{2n+3} = \lim_{n \rightarrow \infty} \frac{1+\frac{1}{n}}{2+\frac{3}{n}} = \frac{1}{2} < 1 \text{ so the series converges by the root test.}$$

(9a). Conditionally convergent: The series converges by the alternating series test since

$$\frac{10}{7n+2} > \frac{10}{7(n+1)+2} \text{ and } \lim_{n \rightarrow \infty} \frac{10}{7n+2} = 0 \text{ but not absolutely since } \sum_{n=1}^{\infty} \left| (-1)^n \frac{10}{7n+2} \right| = \sum_{n=1}^{\infty} \frac{10}{7n+2}$$

diverges by comparing it with the harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$, which diverges, using the limit comparison

$$\text{test: } \lim_{n \rightarrow \infty} \frac{10}{7n+2} / \frac{1}{n} = \lim_{n \rightarrow \infty} \frac{10n}{7n+2} = \frac{10}{7} < \infty.$$

(9b). Absolutely convergent: $\sum_{n=1}^{\infty} \left| (-1)^n \frac{1}{\sqrt{n^5}} \right| = \sum_{n=1}^{\infty} \frac{1}{n^{5/2}}$ a convergent p-series with $p = 5/2 > 1$.

(9c). Absolutely convergent: $\sum_{n=0}^{\infty} |(-1)^n 5^{-n}| = \sum_{n=0}^{\infty} 5^{-n}$ is a convergent geometric series with common ratio $r = 1/5 < 1$.

(9d). $\lim_{n \rightarrow \infty} \frac{n^2 - n - 1}{2n^2 + n + 1} = \lim_{n \rightarrow \infty} \frac{1 - \frac{1}{n} - \frac{1}{n^2}}{2 + \frac{1}{n} + \frac{1}{n^2}} = \frac{1}{2} > 0$ so the series diverges by the nth term test for divergence.

(10a). The power series converges when $|x - 1| < 1$ by the ratio test, which gives a radius of convergence 1 and interval of convergence centered at 1. The series diverges at $x = 2$ (harmonic series) but converges at $x = 0$ (alternate harmonic series), so the interval of convergence is $0 \leq x < 2$.

(10b). The power series converges when $|x - 1| < 1$ by the ratio test, which gives a radius of convergence 1 and interval of convergence centered at 1. The series diverges at $x = 0$ (harmonic series) but converges at $x = 2$ (alternate harmonic series), so the interval of convergence is $0 < x \leq 2$.

(10c). The power series converges when $|x + 1| < 5$ by the ratio test, which gives a radius of convergence 5 and interval of convergence centered at -1 . The series diverges at $x = 4$ (harmonic series) but converges at $x = -6$ (alternate harmonic series), so the interval of convergence is $-6 \leq x < 4$.

(10d). The power series converges when $|x + 1| < 5$ by the ratio test, which gives a radius of convergence 5 and interval of convergence centered at -1 . The series diverges at $x = -6$ (harmonic series) but converges at $x = 4$ (alternate harmonic series), so the interval of convergence is $-6 < x \leq 4$.

$$(11a). \quad p_2(x) = e^2 - 2e^2(x + 1) + 2e^2(x + 1)^2$$

$$(11b). \quad p_2(x) = 1 - \frac{25}{2} (x - 2\pi)^2$$

$$(12a). \quad p_3(x) = 1 + e - e(x + 1) + \frac{e}{2}(x + 1)^2 - \frac{e}{6}(x + 1)^3$$

$$= 1 + \frac{e}{3} - \frac{e}{2}x - \frac{e}{6}x^3$$

$$(12b). \quad p_3(x) = 1 - \frac{1}{2} (x - \frac{\pi}{2})^2$$